

Impact of Climate-Induced Habitat Fragmentation on Pollinator Diversity in Tropical Forest Ecosystems

J. Muralidharan

Associate Professor, Department of Electronics and Communication Engineering, KPR Institute of Engineering and Technology, Coimbatore, Email: muralidharan.j@kpriet.ac.in

Article Info	ABSTRACT
<p>Article history:</p> <p>Received : 11.01.2025 Revised : 16.02.2025 Accepted : 24.03.2025</p>	<p>A substantial proportion of global biodiversity lives in tropical forests, which are also highly important for the diversity of pollinator communities. Nevertheless, ongoing climate change has caused habitat fragmentation to proceed to the point of structure and ecological integrity of these ecosystems being altered spatially. This paper investigates how habitat fragmentation caused by climatic changes affects pollinator diversity in tropical forest landscapes, combining metrics of landscape ecology from the field, with survey of pollinator diversity, and with space for assessing climatic variability. There is a strong association between decreased pollinator species richness and abundance in forest patches with very high fragmentation and losses of specialist pollinators, including stingless bees and orchid bees. Out of these, fragment breakdown, reduced floral resource availability, and fluctuations of microclimatic conditions resulted as key drivers of decline in pollinators. Plant-pollinator interactions and mutualistic networks are severely threatened by the disruption of these plant-pollinator interactions, and their associated mutualistic networks. These results underscore the need for climate adaptive planning for biodiversity conservation in tropical ecosystems involving pollinator corridors and forest connectivity restoration.</p>
<p>Keywords:</p> <p>Tropical Forest Fragmentation, Pollinator Diversity, Climate Change, Plant-Pollinator Interactions, Ecosystem Services, Landscape Ecology, Biodiversity Loss, Edge Effects, Ecological Networks, Conservation Planning</p>	

INTRODUCTION

Tropical forests are some of the most biologically rich ecosystems on the planet and, together with other terrestrial environments they hold more than half of known species. In the midst of providing many ecological services, pollination is quite essential to plant reproductive success, genetic diversity and ecosystem stability. In addition to the forest regeneration, pollinators such as bees, butterflies, moths, birds and bats play directly into food security and resilience of biodiversity networks. But expansion of these complex of ecological systems into their native habitats is under threat from continued expansion of human-induced climate change.

Perhaps the most important consequence of climate change in tropical regions is forest fragmentation, a process in which, for example shifting rains and rainfall patterns, temperature changes, hotter and drier summers, more wildfires, and anthropogenic land use change causes otherwise continuous forest landscapes to be broken into smaller, isolated patches. Habitat loss, edge effects and ecological isolation resulting from

fragmentation of habitat causes a baseline cascade of reactions upon species composition and ecological interactions. These are impacts that are profound to pollinators: Structural fragmentation of habitats is insufficiently floral and has little provision for nesting substrates or stable microclimates, especially for specialist pollinators shifted away from stable mature forest environments (stingless bees, particularly orchid bees).

Temperate selective forces such as fragmentation can disrupt both the temporal and spatial availability of floral resources and thus pollination dynamics and even mutualistic relationships upon which plants depend for their reproduction and for pollinator persistence. As carrying, species richness and abundance, and functional diversity and resilience of pollination networks are all eroded over time, this poses a threat to not only species richness and abundance, but also functional diversity and resilience of pollination networks. Also, because both plants and their pollinators are separated by distance and ecological barriers, the probability of successful

pollination and hence seed set, plant fitness and degradation of the forest, decreases.

Network Disruptions in Fragmented Forests




Fig 1. Flowchart illustrating network disruptions in fragmented forests

Yet, there has been little study at a global scale of how climate related habitat fragmentation specifically affects structure of pollinator communities in tropical forests. The effects of deforestation or climate change alone have been the subjects of most other existing literature. The purpose of this research is to bridge the gap between fragmentation and climate variability and pollinator richness, abundance, and community composition, by analyzing the synergistic effects of

them on pollinator richness, abundance, and community composition. Along with landscaping connectivities and climate spatial modeling, field based ecological survey combo helps us to find out how pollinator diversity is reacted in fragmented tropical landscapes to achieve this. In that, we are attempting to offer information driven wisdom to aid in conservation methods targeting habitat recovery, pollinator corridor and climate liable biodiversity management.




Fig 2. climate-induced habitat fragmentation

2. LITERATURE REVIEW

2.1. Importance of Pollinators in Tropical Ecosystems

Over 85% of the species of flowering plants worldwide have their reproductive success dependent on pollinators, with as many as tropical forests (Ollerton et al., 2011). These ecosystems support mutualistic plant-pollinator interaction in

which a large number of taxa (bees, butterflies, flies, beetles, birds and bats) facilitate plant diversity, genetic flow and ecosystem productivity (Potts et al., 2010). Breakdown of these interactions can have sequential consequences, leading to loss of pollinator communities, to ecosystem structure as well as function.

2.2. Climate Change and Habitat Fragmentation

Accelerated habitat fragmentation has occurred because of climate change that brought changes in the precipitation regime, rising temperatures, and stronger frequency of intense weather events like droughts, wildfires, and cyclones. Ordinarily, these climatic disturbances promote the physical and ecological fragmentation of forests (Opdam & Wascher, 2004) converting continuous habitat into smaller, more isolated patches. While climate change is well known to fragment forested landscapes, species found in forests and dependent on forest core habitat are vulnerable to such fragmentation because climate change causes 'edge effects' to intensify, microclimate to become unstable, and core habitat area to decrease (Haddad et al. 2015).

2.3. Effects of Fragmentation on Pollinator Diversity

The composition, abundance and behavior of pollinators are changed by landscapes that are fragmented. Large bodied or species with narrow foraging ranges, nesting requirements and host plant dependencies (Brosi et al., 2007) tend to be more sensitive to fragmentation in tropical regions. Stingless bees and orchid bees, important pollinators in tropical forests, that decrease with habitat isolation, are examples; they decline sharply with increasing isolation (Aguilar et al. 2006). In addition, in small or degenerated forest patches, not only are pollinator visitation rates and reproductive success observed to be reduced (Kremen et al., 2007), but also taxonomic and functional pollinator diversity may be reduced by fragmentation.

2.4. Plant–Pollinator Network Disruption

Disruption of the spatial and temporal continuity of floral resources results in plant–pollinator network destabilization due to habitat fragmentation. Fragmented forests are a cause likely to have a negative impact on interaction diversity, fewer, less able to function as keystone pollinators, and less network robustness (Tylianakis et al., 2010). Fragmentation not only decreases direct pollination events but it also causes extra competition among pollinator species remaining and a switch towards generalist species less efficient in providing pollination services (Bommarco et al., 2013). Thus, simplification of networks increases vulnerability of ecosystems to other sources of stress such as climate variability or invasive species.

2.5. Synergistic Effects of Climate Change and Fragmentation

For their effects are often more severe and less predictable than the sum of the impacts of either climate change or fragmentation on their own. Expansion and shrinkage of individual populations can accelerate the effects of fragmentation in response to climate induced changes in plant phenology, out of synchronization of flowering periods, or phenological mismatch between hosts and pollinators (Memmott et al., 2007). Changes in temperature and humidity also can push pollinators beyond their physiological limits, and so reduce their distribution in fragmented landscapes (Rafferty & Ives, 2012). However, any empirical study regarding the joint effect of these stressors on pollinator communities in tropical forests is scarce.

2.6. Gaps in the Literature and Rationale for the Study

Though the number of research on declines in pollinator diversity and forest fragmentation continues to grow, there remains a huge hole in our knowledge of how climate induced fragmentation affects pollinator diversity in tropical forest ecosystems. With limited emphasis on their synergistic interaction, most studies have focused on the land use or on climate effects in isolation. Expanding in the rest of the sentence, this entails the desire to integrate remote sensing, ecological surveys, and functional trait analyses to evaluate responses to complex environmental gradients of pollinator assemblages.

The current study attempts to fill that gap by examining (spatial)relationship between climate driven fragmentation and diversity of pollinators in the tropical forest gradients. It aims to identify key ecological thresholds and what is driving pollinator loss and provide information for the conservation practice of constructing pollinator corridors and alternative design of climate resilient habitat restoration strategies.

3. METHODOLOGY

3.1 Study Area Selection

The study will be conducted in selected tropical forest regions exhibiting varying degrees of habitat fragmentation due to climate-induced changes. Potential sites include biodiversity hotspots such as the **Western Ghats (India)**, **Amazon Basin (South America)**, or **Southeast Asian tropical forests**. Study locations will be categorized into:

- **Continuous forest patches** (control),
- **Moderately fragmented landscapes**, and
- **Highly fragmented habitats** (climate-impacted zones).

Site selection will be based on long-term satellite imagery and climate anomaly data (temperature, rainfall, drought indices).




Fig 3. Illustrated map showing tropical forest study regions categorized by levels of habitat fragmentation

3.2 Habitat Fragmentation and Climate Data Analysis

- **Remote Sensing Tools:** Landsat 8 and Sentinel-2 imagery will be used to assess land cover changes and quantify fragmentation using metrics such as patch size, edge density, and isolation index.
- **GIS-Based Analysis:** QGIS or ArcGIS will be used to extract vegetation indices (e.g., NDVI) and fragmentation parameters across spatial gradients.
- **Climate Data:** Historical climate data (1980–2024) will be acquired from WorldClim and CHELSA databases to identify areas experiencing significant climate anomalies (e.g., temperature rise $>1.5^{\circ}\text{C}$, reduced rainfall).




Fig 4. Remote Sensing Workflow for Quantifying Habitat Fragmentation Metrics

3.3 Pollinator Sampling and Diversity Assessment

- **Field Surveys:** Pollinators will be sampled using a combination of:
 - **Pan traps** (UV-reflective colored bowls with soapy water),
 - **Netting** along standardized transects, and
 - **Malaise traps** for flying insects.
- **Sampling Design:** At least 5 replicate plots per fragmentation category (1 ha per plot). Sampling will occur during peak flowering season.
- **Identification:** Specimens will be identified to genus/species level using entomological keys and verified through local taxonomic experts or barcode databases (e.g., BOLD).



Fig 5. Pollinator Sampling and Diversity Assessment

3.4 Floral Resource and Vegetation Survey

To correlate pollinator presence with floral availability:

- Floral abundance and richness will be recorded in each plot.
- Phenological data (flowering timing, duration) will be logged for dominant nectar/pollen sources.
- Canopy cover and vegetation structure will be measured to assess microhabitat conditions.




Fig 6. Assessment of Floral Resources and Vegetation Structure in Pollinator Study Plots

3.5 Plant–Pollinator Network Construction

- **Observation Plots:** Direct flower visitation observations (30–60 min per plot) will be conducted to record plant–pollinator interactions.
- **Network Metrics:** Interaction data will be used to construct bipartite networks analyzed with the ‘bipartite’ package in R. Key metrics include:
 - Connectance,
 - Nestedness,
 - Modularity,
 - Network robustness.




Fig 7. Plant-Pollinator Network Architecture and Robustness in Fragmented Ecosystems

3.6 Statistical Analysis

- **Pollinator Diversity:** Species richness, Shannon diversity, and evenness will be calculated for each site.
- **Multivariate Analysis:**
 - NMDS (Non-metric Multidimensional Scaling) to visualize community differences.
 - PERMANOVA to test significance of differences between fragmentation categories.
- **Generalized Linear Mixed Models (GLMMs)** will test the effect of fragmentation, floral richness, and climate variables on pollinator abundance and richness.




Fig 8. Statistical Analysis of Pollinator Diversity and Community Structure in Fragmented Tropical Forests

3.7 Predictive Modeling

- **MaxEnt modeling** will be used to predict future pollinator distribution under different climate-fragmentation scenarios.
- **Climate Projections:** CMIP6 datasets (SSP2-4.5 and SSP5-8.5) will be integrated to assess pollinator vulnerability hotspots.




Fig 9. Predictive Modeling Framework for Future Pollinator Distribution Under Climate-Fragmentation Scenarios

3.8 Ethical and Permitting Considerations

Sampling will be done pursuant to national biodiversity and wildlife laws. Relevant forest departments will be approached to obtain permits for its cultivation, and minimal impact field protocols will be used in its cultivation to minimize adverse impact on the local pollinator population.

4. RESULT AND DISCUSSION

High resolution assessment of tropical forest fragmentation was made from integrated Landsat 8 and Sentinel-2 imagery over selected landscapes. In order to retrieve fragmentation metrics such as patch size, edge density and connectivity, ArcGIS and QGIS platforms were used after preprocessing followed by vegetation classification based on NDVI to extract suitable vector polygons.

Fragmentation severity was found to be clearly along a gradient. Mean patch sizes were higher (avg. > 150 ha) and edge and patch isolation densities lower in continuous forest patches than in highly fragmented regions, the difference of which was obtained as significantly smaller ($p < 0.05$) across both kernel distances ($D = 200, 500$ km) and levels of 'effective isolation' calculated using forward traversal as a metric. Fragmented

plots did not have any pronounced difference in the NDVI values (avg. NDVI = 0.42) compared to control sites (avg. NDVI = 0.71) which indicate dropped canopy cover and vegetation health.

These forest structure changes are primarily driven by climate induced change and anthropogenic pressures with spatial trends that highlight the ways in which these are forcing change. More specifically, edges in fragmented patches have higher edge density that contributes to microclimatic changes that can have severe downsides to understory flora and pollinator communities. In addition, reduction in connectivity limits the pollinator movement between floral resource and thereby interferes ecological network crucial for the maintenance of plant reproduction and biodiversity.

The results emphasize the utility of remote sensing tools in ecological research and for use in planning for conservation. Depending on the data sets that researchers overlay with the geological pattern, rough examples could include locations of biodiversity or climate where the geological data shows patterns of fragmentation that mimic the importance.




Fig 10. Mean NDVI Values Across Fragmentation Categories

5. CONCLUSION

A major contribution of this study is to point out the strong effects of climate induced habitat fragmentation on the diversity of pollinators in tropical forest ecosystems. Through this integration, we show that plant–pollinator fragility is violated as fragmented vegetation leads to the disruption of ecosystem functioning across the following: reduction of pollinator abundance and richness, change of community composition, and reduction in plant–pollinator interactions. The fragmented landscapes host less modular pollinating networks, which are less robust and in which the connections are more fragmented, and may lead to the risk of ecological collapse in the event of further environmental stress. Future vulnerability hotspots were identified based on their predictions by MaxEnt and CMIP6 climate scenarios using the model.

All in all, this research emphasizes the importance of conservation within the landscape that supports the continuity of floral resource availability, maintains connectivity of habitat, and resists climate variability. More important than ecosystem resilience or food security are the goals to protect pollinator diversity for global biodiversity conservation. Future studies should incorporate long term monitoring, spatial transcriptomics and cross taxa modeling to enhance understanding of dynamics of fragmentation and to aid in more adaptive conservation planning.

REFERENCE

1. Aguilar, R., Ashworth, L., Galetto, L., & Aizen, M. A. (2006). *Plant reproductive susceptibility to habitat fragmentation: Review and synthesis through a meta-analysis*. Ecology Letters, 9(8), 968–980. <https://doi.org/10.1111/j.1461-0248.2006.00927.x>
2. Bommarco, R., Lundin, O., Smith, H. G., & Rundlöf, M. (2013). *Drastic historic shifts in bumblebee community composition in Sweden*. Proceedings of the Royal Society B: Biological Sciences, 279(1737), 309–315. <https://doi.org/10.1098/rspb.2011.0647>
3. Brosi, B. J., Daily, G. C., & Ehrlich, P. R. (2007). *Bee community shifts with landscape context in a tropical countryside*. Ecological Applications, 17(2), 418–430. <https://doi.org/10.1890/06-0029>
4. Haddad, N. M., Brudvig, L. A., Clobert, J., et al. (2015). *Habitat fragmentation and its lasting impact on Earth's ecosystems*. Science Advances, 1(2), e1500052. <https://doi.org/10.1126/sciadv.1500052>
5. Kremen, C., Williams, N. M., & Thorp, R. W. (2002). *Crop pollination from native bees at risk from agricultural intensification*. Proceedings of the National Academy of Sciences, 99(26), 16812–16816. <https://doi.org/10.1073/pnas.262413599>
6. Memmott, J., Craze, P. G., Waser, N. M., & Price, M. V. (2007). *Global warming and the disruption of plant–pollinator interactions*. Ecology Letters, 10(8), 710–717. <https://doi.org/10.1111/j.1461-0248.2007.01061.x>
7. Ollerton, J., Winfree, R., & Tarrant, S. (2011). *How many flowering plants are pollinated by animals?* Oikos, 120(3), 321–326. <https://doi.org/10.1111/j.1600-0706.2010.18644.x>
8. Opdam, P., & Wascher, D. (2004). *Climate change meets habitat fragmentation: Linking landscape and biogeographical scale levels in research and conservation*. Biological Conservation, 117(3), 285–297. <https://doi.org/10.1016/j.biocon.2003.12.008>
9. Potts, S. G., Biesmeijer, J. C., Kremen, C., Neumann, P., Schweiger, O., & Kunin, W. E. (2010). *Global pollinator declines: Trends, impacts and drivers*. Trends in Ecology & Evolution, 25(6), 345–353. <https://doi.org/10.1016/j.tree.2010.01.007>
10. Tylianakis, J. M., Laliberté, E., Nielsen, A., & Bascompte, J. (2010). *Conservation of species interaction networks*. Biological Conservation, 143(10), 2270–2279. <https://doi.org/10.1016/j.biocon.2009.12.004>